

CHIBA-EP-70

February, 1994

hep-th/9404026

Supersymmetry in Stochastic Quantization Method and Field-Dependent Kernel

KENJI IKEGAMI

*Graduate School of Science and Technology,
Chiba University,
1-33 Yayoi-cho, Inage-ku, Chiba 263, JAPAN*

ABSTRACT

We define a discretized Langevin equation in Stratonovich-*type* calculus. We show that a generating functional with a field-dependent kernel can be written in mid-point prescription only when we calculate in the calculus. Moreover we investigate whether supersymmetry of the stochastic action with field-dependent kernel exists or not.

1. INTRODUCTION

The stochastic quantization method (SQM) was first proposed by Parisi and Wu as an alternative quantization method in 1981.^{[1][2]} SQM can be applied to gauge theories without the gauge fixing procedure, i.e. without Faddeev-Popov ghost fields. Instead of introducing ghost field, the method produces the same contribution as the path-integral quantization method (PIQM). This fact was already confirmed perturbatively for Yang-Mills fields^[3] and for non-Abelian anti-symmetric tensor fields.^[4]

SQM has a powerful tool, “kernel”, which, among others, gives new regularization schemes.^[5] Kernel is also introduced for systems including massless fermion.^[6] Moreover, “field-dependent” kernel is introduced for systems including graviton,^[7] systems with spontaneously broken symmetry,^[8] and bottomless systems.^[9] On the other hand, it is well known that theories quantized stochastically display supersymmetry.^{[10][11]} So my question is whether SQM with *field-dependent* kernel has supersymmetry or not. In this paper we show that SQM with field-dependent kernel has supersymmetry as well as the one without kernel. While Ref.[12] showed that stochastic action with field-dependent kernel was invariant by operation with the supersymmetry generator[‡] Q , it did not show that the action was invariant by operation with $\overline{Q}$ and could be described in superfield formalism.^[13] Besides, boundary condition^[11] of generating functionals function was not discussed in Ref.[12].

When we construct the stochastic action, Ito^[14] or Stratonovich^[15] calculus can be used. Leibnitz rule, which is indispensable to supersymmetry, can be used in the Stratonovich calculus, but cannot be used in Ito calculus in continuum limit. Therefore, we define a discretized Langevin equation in Stratonovich-*type* calculus, and we construct the stochastic action where Leibnitz rule can be used. In this paper, we show that only in the calculus the stochastic action in *mid-point prescription* can be constructed.

‡ The definition of $Q, \overline{Q}$ are given in this paper

2. BOUNDARY CONDITION OF GENERATING FUNCTIONAL

First, we take up a system with variables $q(x)$ and the classical action $S(q)$ in n -dimensional space-time. To quantize the system stochastically, we consider the fictitious time interval $[-T, T]$ and divide the interval to $2N$ segments with space T/N . We shall let T tend to infinity later. Besides, we define the discretized Langevin equation with the field-dependent kernel $K(\bar{q}_i)$

$$\begin{aligned} dq_i(x) &\equiv q_{i+1} - q_i = -K(\bar{q}_i) \frac{\delta S}{\delta \bar{q}_i} + R_i \frac{\delta R_i}{\delta \bar{q}_i} + R_i dW_i, \\ i &= 1, 2, \dots, N, \quad -T = t_{-N} < t_{-N+1} < \dots < t_N = T, \\ q_i(x) &\equiv q(x, t_i), \quad \bar{q}_i \equiv \frac{1}{2}(q_{i+1} + q_i), \quad R_i \equiv R(\bar{q}_i) \\ K(\bar{q}_i) &= R^2(\bar{q}_i), \quad dW_i(x) \equiv W(x, t_{i+1}) - W(x, t_i), \end{aligned} \tag{1}$$

where $dt = T/N$ and we assume the kernel $K(\bar{q}_i)$ is positive-definite. $W(x, t_i)$ is a Wiener process defined by the following correlation

$$\begin{aligned} \langle (W(x, t_i) - W(x, t_k))(W(y, t_j) - W(y, t_l)) \rangle &= 2(t_j - t_k) \delta^n(x - y), \\ i &> j > k > l, \\ \langle \dots \rangle &= \int D(dW_i) (\dots) \exp\left[-\frac{1}{4} \int d^n x \sum_{i=-N}^N \left(\frac{dW_i}{dt}\right)^2 dt\right]. \end{aligned} \tag{2}$$

The expression of eq.(1) is different from the ordinary one in Stratonovich calculus^[17] or the one in Stratonovich-related calculus,^[18] but we want to regard it as the expression in Stratonovich-*type* calculus. The advantage of the calculus is shown later.

Now we are able to show that Leibnitz rule can be used in the Stratonovich-type calculus in the continuum limit $dt \rightarrow 0$ as follows. We shall calculate the expectation value

$$\begin{aligned} &\int d^n y \left\langle \frac{\delta f(\bar{q}_i)}{\delta \bar{q}_i(y)} dq_i(y) \right\rangle \\ &= \int d^n y \left\langle \frac{\delta f(\bar{q}_i)}{\delta \bar{q}_i(y)} \left\{ (-K(\bar{q}_i) \frac{\delta S(\bar{q}_i)}{\delta \bar{q}_i(y)} + R_i \frac{\delta R_i}{\delta \bar{q}_i(y)}) dt + R_i dW_i(y) \right\} \right\rangle, \\ &= \int d^n y \left\langle \frac{\delta f(\bar{q}_i)}{\delta \bar{q}_i(y)} \left(-K(\bar{q}_i) \frac{\delta S(\bar{q}_i)}{\delta \bar{q}_i(y)} + 2R_i \frac{\delta R_i}{\delta \bar{q}_i(y)} \right) dt + K(\bar{q}_i) \frac{\delta^2 f(\bar{q}_i)}{\delta \bar{q}_i(y)^2} dt \right\rangle, \end{aligned} \tag{3}$$

where $f(\bar{q}_i)$ is some function of $\bar{q}_i$. In second equality we use the fact that $\langle q_i dW_i \rangle = 0$, $\langle q_{i+1} dW_i \rangle = \langle 2R(\bar{q}_i)dt \rangle + O(dt^2)$. The final expression in eq.(3) is equivalent to the value

$$\int d^n y \langle \frac{\delta f(q_i)}{\delta q_i(y)} (-K(q_i) \frac{\delta S(q_i)}{\delta q_i(y)} + 2R(q_i) \frac{\delta R(q_i)}{\delta q_i(y)}) dt + K(q_i) \frac{\delta^2 f(q_i)}{\delta q_i^2(y)} dt \rangle, \quad (4)$$

up to dt because $\bar{q}_i = q_i + \frac{1}{2}dq_i$. The expression coincides with $\langle df(q_i) \rangle \equiv \langle f(q_{i+1}) - f(q_i) \rangle$ in Ito calculus because the Langevin equation in Ito calculus is

$$dq_i = -K(q_i) \frac{\delta S(q_i)}{\delta q_i} dt + 2R(q_i) \frac{\delta R(q_i)}{\delta q_i} dt + R(q_i) dW_i. \quad (5)$$

Eqs.(3) and (4) mean

$$\langle df(q_i) \rangle = \int d^n y \langle \frac{\delta f(\bar{q}_i)}{\delta \bar{q}_i(y)} dq_i(y) \rangle + O(dt^2). \quad (6)$$

Therefore, Leibnitz rule can be used in continuum limit of the Stratonovich-type calculus.

Next, let us introduce the generating functional

$$\begin{aligned} Z[J] &\equiv \int D(dW_i) \exp \left[- \int d^n x \sum_{i=-N}^N \left\{ \frac{1}{4} \left(\frac{dW_i}{dt} \right)^2 + J_i(x) q_i^W(x) \right\} dt \right], \\ &= \int Dq_i \det(M_{ij}) \\ &\times \exp \left[- \int d^n x \sum_{i=-N}^N \left[\frac{1}{4} \left\{ \left(\frac{dq_i}{dt} + K \frac{\delta S(q_i)}{\delta \bar{q}_i} - R_i \frac{\delta R_i}{\delta \bar{q}_i} \right) R_i^{-1} \right\}^2 + J_i(x) q_i^W(x) \right] dt \right], \\ M_{ij} &\equiv \frac{\delta(dW_i)}{\delta q_{j+1}}, \end{aligned} \quad (7)$$

where q_i^W is a solution of eq.(1). Another expression of the generating functional

is

$$\begin{aligned}\tilde{Z}[J] &= \int (Dq Dp D\bar{C} DC) e^{-A + \int d^n x dt J(x,t) q(x,t)}, \\ A &\equiv \int d^n x \int_{-T}^T dt \{ p K(q) p - i p (\dot{q} + K \frac{\delta S}{\delta q} - R \frac{\delta R}{\delta q}) \\ &\quad - \bar{C} R(q) \frac{\partial}{\partial q} R^{-1} (\dot{q} + K \frac{\delta S}{\delta q} - R \frac{\delta R}{\delta q}) C \},\end{aligned}\tag{8}$$

in continuum limit. Here the auxiliary field p and Grassmannian variables $\bar{C}, C$ are introduced. In order to assert that $Z = \tilde{Z}$, we need to prove

$$\lim_{dt \rightarrow 0} \det(M_{ij}) = I_{fermi},\tag{9}$$

where

$$\begin{aligned}I_{fermi} &\equiv \int Dp DC D\bar{C} e^{-A_f}, \\ A_f &\equiv \int d^n x \int_{-T}^T dt \{ (p - \frac{i}{2} (\dot{q} + K \frac{\delta S}{\delta q} - R \frac{\delta R}{\delta q}) K^{-1}) R \}^2 \\ &\quad - \bar{C} R(q) \frac{\partial}{\partial q} R^{-1} (\dot{q} + K \frac{\delta S}{\delta q} - R \frac{\delta R}{\delta q}) C \}.\end{aligned}\tag{10}$$

In order to calculate $\det(M_{ij})$ and I_{fermi} , we take the twisted boundary condition^[11]

$$q(-T) = e^{-i\nu} q(T), \quad C(-T) = e^{-i\nu} C(T), \quad e^{-i\nu} \bar{C}(-T) = \bar{C}(T), \quad e^{-i\nu} p(-T) = p(T).\tag{11}$$

The boundary condition for $\nu = 0$ corresponds to the periodic one

$$q(-T) = q(T), \quad C(-T) = C(T), \quad \bar{C}(-T) = \bar{C}(T), \quad p(-T) = p(T),\tag{12}$$

or for $\nu = -i\infty$ the causal and anti-causal one

$$q(-T) = 0, \quad C(-T) = 0, \quad \bar{C}(T) = 0, \quad p(T) = 0,\tag{13}$$

In this paper, we choose ν as $Re(\nu) = 0$ because q is a real variable. We can

calculate $\det(M_{ij})$ as

$$\begin{aligned} \det(M_{ij}) &= \prod_{i=-N}^N R_i^{-1} \\ &\times \det \begin{pmatrix} 1 + G_{-N}dt & & e^{-i\nu}(-1 + G_{-N}dt) \\ -1 + G_{-N+1}dt & 1 + G_{-N+1}dt & \\ & & -1 + G_Ndt & 1 + G_Ndt \end{pmatrix}, \\ G_i dt &\equiv \frac{1}{2} R_i \frac{\delta R_i^{-1}}{\delta \bar{q}_i} (q_{i+1} - q_i) + \frac{1}{2} R_i \frac{\partial}{\partial \bar{q}_i} \left(R_i \frac{\delta S(\bar{q}_i)}{\delta \bar{q}_i} - \frac{\delta R_i}{\delta \bar{q}_i} \right) dt. \end{aligned} \quad (14)$$

The Stratonovich-*type* calculus has the advantage that the $\det(M_{ij})$ is expressed in the mid-point prescription, i.e.,

$$\det(M_{ij}) = \prod_{i=-N}^N g\left(\frac{q_{i+1} + q_i}{2}, q_{i+1} - q_i\right), \quad (15)$$

where g is some function. In ordinary Stratonovich calculus, $\det(M_{ij})$ cannot be expressed in the prescription in case that kernel is field-dependent. By straightforward calculation, the determinant is

$$\begin{aligned} \lim_{dt \rightarrow 0} \det(M_{ij}) &= \prod_{t=-T}^T R^{-1}(t) \{e^{\int_{-T}^T dt d^n x G(t)} - e^{-\int_{-T}^T dt d^n x G(t) - i\nu}\}, \\ &= \prod_t R^{-1}(t) \sinh\left\{\int_{-T}^T dt d^n x G(t) + \frac{1}{2}i\nu\right\} \times \text{Constant}. \end{aligned} \quad (16)$$

Next I_{fermi} can be also calculated as

$$I_{fermi} = \sinh\left[\left\{\frac{1}{2}i\nu + \int_{-T}^T d^n x dt G(t)\right\}\right]. \quad (17)$$

by following ref.[16]. From eqs.(16) and (17), we have proved that Z is equivalent to $\tilde{Z}$ when the boundary condition (11) is taken.

3. SUPERSYMMETRY AND FIELD-DEPENDENT KERNEL

From now on, we introduce the field-dependent kernel again. The expression of A in (8) is rather complicated and it is difficult to recognize whether the stochastic action with field-dependent kernel has supersymmetry or not. In fact, A is not invariant under the supersymmetric transformations under which the stochastic action without kernel is invariant.

Here, we consider the change of variables

$$q' = \int dq R^{-1}(q), \quad p' = R(q)p, \quad \overline{C}' = \overline{C}R(q), \quad C' = R^{-1}(q)C. \quad (18)$$

This leads to

$$\begin{aligned} \tilde{Z}[J] &= \int Dp' Dq' DC' D\overline{C}' e^{-A' + \int d^n x dt J(x,t)q(q')} \\ A' &\equiv \int d^n x \int_{-T}^T dt \left\{ p'^2 - ip'(\dot{q}' + \frac{\delta S}{\delta q'} - R^{-1} \frac{\delta R}{\delta q'}) - \overline{C}' \frac{\partial}{\partial q'} (\dot{q}' + \frac{\delta S}{\delta q'} - R^{-1} \frac{\delta R}{\delta q'}) C' \right\}, \end{aligned} \quad (19)$$

where we assume that the first relation in eq.(18) can be solved for q in terms of q' . The periodic boundary condition (12) can be also expressed in terms of new variables as

$$q'(-T) = q'(T), \quad C'(-T) = C'(T), \quad \overline{C}'(-T) = \overline{C}'(T), \quad p'(-T) = p'(T), \quad (20)$$

i.e., periodic boundary condition. For $\nu = -i\infty$, the boundary condition (13) can be also expressed as

$$q'(-T) = 0, \quad C'(-T) = 0, \quad \overline{C}'(T) = 0, \quad p'(T) = 0 \quad (21)$$

i.e., causal and anti-causal boundary condition, where we choose the integral constant as $\int R^{-1} dq|_{q=0} = 0$.

The stochastic action A' with the boundary condition (20) is invariant under the super-transformations

$$\delta q' = \bar{\epsilon} C', \quad \delta \bar{C}' = -i\bar{\epsilon} p', \quad \delta C' = 0, \quad \delta p' = 0, \quad (22)$$

and

$$\delta q' = \bar{C}' \epsilon, \quad \delta \bar{C}' = 0, \quad \delta C' = -i\epsilon p' - \epsilon \dot{q}', \quad \delta p' = i\dot{\bar{C}}' \epsilon. \quad (23)$$

On the other hand, A' with (21) is not invariant under the transformation (23) as well as the case of no kernel.^[11]

In terms of original variables $q, C, \bar{C}$ and p , the transformation (22) is expressed as

$$\delta q = \bar{\epsilon} C, \quad \delta \bar{C} = \bar{\epsilon} \bar{C} \frac{\partial R}{\partial q} R^{-1} C - i\bar{\epsilon} p, \quad \delta C = -R \frac{\partial R^{-1}}{\partial q} \bar{\epsilon} C, \quad \delta p = -R^{-1} \frac{\partial R}{\partial q} \bar{\epsilon} C p, \quad (24)$$

and the transformation (23) is

$$\begin{aligned} \delta q &= \bar{C} K(q) \epsilon, \quad \delta \bar{C} = 0, \\ \delta C &= -i\epsilon K(q) p - \epsilon \dot{q} + \bar{C} \epsilon \frac{\partial R(q)}{\partial q} R(q) C, \\ \delta p &= i\dot{\bar{C}} \epsilon + i\bar{C} \frac{\partial R(q)}{\partial q} R^{-1} \dot{q} \epsilon - \bar{C} \epsilon \frac{\partial R(q)}{\partial q} R(q) p. \end{aligned} \quad (25)$$

Finally, we express the stochastic action in terms of the superfield Φ' as

$$\begin{aligned} A' &= -\int d^n x d^2 \theta \int_{-T}^T dt \{ \bar{D}_\theta \Phi' D_{\bar{\theta}} \Phi' + L(q(\Phi')) - \delta^n(0) \ln R(q(\Phi')) \}, \\ D_{\bar{\theta}} &\equiv \partial_{\bar{\theta}} - \theta \partial_t, \quad \bar{D}_\theta \equiv \partial_\theta, \quad \Phi' \equiv q' + \bar{\theta} C' + \bar{C}' \theta - i\bar{\theta} \theta p', \end{aligned} \quad (26)$$

where $\theta, \bar{\theta}$ are Grassmannian variables. As discussed above, the expression is invariant under operation with $Q(\equiv \partial_{\bar{\theta}})$ and $\bar{Q}(\equiv \bar{\partial}_\theta + \bar{\theta} \partial_t)$ for $\nu = 0$ and invariant under the operation with only Q for $\nu = -i\infty$

4. SUMMARY

We defined the discretized Langevin equation (1) with field-dependent kernel in Stratonovich-*type* calculus in which Leibnitz rule can be used and constructed the generating functional. When the field-dependent kernel is introduced, only the generating functional constructed in the Stratonovich-*type* calculus can be expressed in the prescription. Besides, we showed that the generating functional (7) constructed from Wiener process distribution is equivalent to the continuous one (8) when we take the twisted boundary condition (11).

Moreover, we showed that the stochastic action with the periodic boundary condition (12) is invariant under the super-transformations (22) and (23) when field-dependent kernel is introduced. The stochastic action with the causal and anti-causal boundary condition (13) is not invariant under the super-transformation (23) as well as the one without kernel. We also show that the stochastic action can be also described in terms of superfield.

Acknowledgments

Author thanks Prof.S.Kawasaki, Drs.R.Mochizuki and A.Nakamura for valuable discussion and comments.

REFERENCES

1. G.Parisi and Y.Wu, Sci.Sin. **24** (1981)483.
2. For a review, P.H.Damgaard and H.Hüffel, Phys.Rep.**152**(1987)227.
3. M.Namiki, I.Ohba, K.Okano and Y.Yamanaka, Prog. Theor. Phys. **69** (1983)1580.
4. A.Nakamura, Prog.Theor.Phys. **86**(1991)925.
5. For example, J.T.Breit, S.Gupta and A.Zaks, Nucl.Phys.B233 (1984)61;
J.Alfaro, Nucl.Phys.B253(1985)464.
6. U.Kakudo, Y.Taguchi, A.Tanaka and K.Yamamoto, Prog.Theor.Phys.**69** (1983)1225;
T.Fukai, H.Nakazato,I.Ohba,K.Okano and Y.Yamanaka, Prog.Theor.Phys.
69(1983)1600.
7. H.Rumpf, Phys.Rev.D33(1986)942.
8. K.Ikegami, R.Mochizuki and K.Yoshida, Prog.Theor.Phys.**89**(1993)197.
9. S.Tanaka, M.Namiki, I.Ohba, M.Mizutani, N.Komoike and M.Kanenaga,
Phys.Lett.B288(1992)129.
10. G.Parisi and N.Sourlas, Phys.Rev.Lett.43(1979)744;
Nucl.Phys.B206(1983)321.
11. H.Nakazato, K.Okano, L.Schülke and Y.Yamanaka, Nucl.Phys.B346(1990)
611;
S.Marculescu, K.Okano and L.Schülke, Nucl.Phys.B349(1991),463.
12. R.Mochizuki, Prog.Theor.Phys. **85**(1991)407.
13. E.Egorian and S.Kalitzin, Phys.Lett.B129(1983)320;
R.Kirschner, Phys.Lett.B139(1984)180;
E.Gozzi, Phys.Lett.B143(1984)183.
14. K.Ito, Proc.Imp.Acad. **20** (1944)519.
15. R.L.Stratonovich, Conditional Markov Processes and Their Application to
the Theory of Optimal Control(Elsevier, NewYork, 1968).

16. E.Gozzi, Phys. Rev. D28 (1983) 1922.
17. For example, M.Namiki, Stochastic Quantization (Springer-Verlag, 1992)
p.209.
18. H.Ezawa and J.R.Klauder, Prog. Theor. Phys. 74(1985)904.